

LABORATORY OF STEM CELL BIOLOGY



Associate Professor
TSUBOUCHI, Tomomi

NIBB Research Fellow: KAMIKAWA, Yasunao
Technical Assistant: YASUI, Naomi

DNA is constantly damaged from both endogenous and exogenous sources. One of the most important challenges for all living organisms is to prevent genome instability that can lead to malfunction of a cell. Our group is interested in the strategies through which cells protect themselves from alterations in the genome. To date, much information has been gained from various model organisms and tissue culture cells, and we are beginning to learn that the choice of genome-maintenance strategies taken by a cell depends on the cell type, cell cycle- and developmental stages. In the Laboratory of Stem Cell Biology, we are currently focusing our attention on the genome maintenance mechanisms of the embryonic stem cells, and their roles during differentiation and reprogramming processes.

I. Self-renewal of Embryonic Stem Cells and Their Genome-Maintenance Mechanisms

Embryonic stem (ES) cells are derived from the blastocyst stage of embryonic development, and are capable of differentiating into all cell types that compose our body (i.e., ES cells are “pluripotent”). Pluripotent cells exist only transiently and are lost as development proceeds. On the other hand, ES cells are capable of proliferating indefinitely when given an appropriate culturing condition. Curiously, ES cells proliferate with truncated gap phases while S (DNA replication) and M (mitosis) phases are as lengthy as other cell types. ES cells also appear to lack some of the mechanisms that ensure genome integrity (i.e., checkpoint mechanisms), the significance of which remains a mystery.

To date, studies on cell cycle regulation in ES cells have not been straightforward compared to that of other cell types, as many commonly used cell-synchronization protocols are ineffective for ES cells. We have established a protocol to synchronize ES cells using centrifugal elutriation (Tsubouchi et al., *Cell*, 2013), which allowed us to investigate specific stages of the ES cell cycle. We are currently aiming to address why and how ES cells maintain such unique cell cycle regulation and how that is interlinked with pluripotency by carrying out side-by-side analyses between ES cells and differentiated populations.

II. Genome Instability during Nuclear Reprogramming

In order to gain a deeper understanding of the relationship between the choice of genome maintenance mechanisms and pluripotency, we are investigating the behavior of factors involved in genome maintenance mechanisms during nuclear reprogramming towards pluripotency.

Specifically, we take advantage of the cell-to-cell fusion approach, in which a target cell is fused to a pluripotent stem

cell to induce pluripotency within a target nucleus (Figure 1). The cell fusion system is a simple, versatile way to induce reprogramming towards another lineage, not limited to pluripotency. Using this system, the first sign of reprogramming can be detected from within a few hours to one day after fusion, allowing us to monitor the initial events of reprogramming after induction.




Figure 1. Reprogramming induced through cellular fusion: a human lymphoblastoid nucleus (blue, with green spots) is induced to undergo nuclear reprogramming towards pluripotency upon fusion with mouse ES cells (green).

Using this system, we previously found that DNA synthesis is an important event for successful reprogramming (Tsubouchi et al., *Cell*, 2013). Recent reports indicate that reprogramming may cause genetic instabilities, some of which are thought to arise as DNA replication errors. We are currently investigating how such genetic instability may arise, and how it is linked to reprogramming-specific events.

III. Future Perspective

While the fundamental mechanisms that maintain genome integrity have been widely studied using various models, the danger a cell might face when altering their cellular identity (through differentiation, reprogramming etc.) is unknown. Recent studies of cancer genome sequencing repeatedly identified mutations in the factors that govern cellular identities, leading us to hypothesize that cells may experience genome instability when their identity is unstable. Our goal is to uncover the nature of such genetic instability and to gain a comprehensive understanding of the mechanisms that maintain genome integrity.

Publication List:

[Original paper]

- Leung, W.-K., Humphries, N., Afshar, N., Argunhan, B., Terentyev, Y., Tsubouchi, T., and Tsubouchi, H. (2015). The synaptonemal complex is assembled by a polySUMOylation-driven feedback mechanism in yeast. *J. Cell Biol.* 211, 785-793.