

LABORATORY OF MORRHODIVERSITY



Associate Professor
KODAMA, Ryuji

The aim of this laboratory is to observe the variety of morphogenetic processes that occur in the course of the ontogenesis of multicellular organisms and to analyze the mechanisms of such processes, mainly by morphological methods. The accumulation of such analyses of the embryogenetic processes of related species is expected to provide an insight into the evolution of morphogenetic processes. This laboratory uses the wings of lepidopteran insects for our morphological studies.

I . Wing morphogenesis

The wings of the lepidopteran insects (butterflies and moths) develop from the wing imaginal disc, which is a hollow sac made of simple epithelium. Due to its simple construction, this tissue is a good material for studying cellular interactions in the course of morphogenesis.

The outline shape of the adult wing is often different from that of the pupal wing. This difference is brought about by the programmed cell death of the marginal area of the pupal wing. The marginal dying area is called "the degeneration region" and the internal area, which develops into the adult wing, is called "the differentiation region".

Cell death in the degeneration region proceeds very rapidly and finishes in a half-day to one-day period in *Pieris rapae* and in several other examined species. It was shown that the dying cells in the degeneration region have characteristics in common with the apop-totic cell death in mammalian cells, such as fragmented and condensed nuclei containing short DNA fragments detected by TUNEL staining. The cells in the degeneration region are actively en-gulfed by the macrophages in the cavity beneath the wing epithelium. The macrophages seem to be concentrated beneath the degeneration region by the strong adhesion between the basal surfaces of the dorsal and ventral epithelium in the differentiation region, which occurs at the time of prominent cell death and excludes macrophages out of the differentiation region. This realized concentration of macrophages seems to accelerate the shrinkage of the degeneration region.

A possible physiological role of the cell degeneration at the wing margin is to make space for the growth of the marginal scales. Marginal scales are extremely elongated scales that grow densely on the edge of the wing. These scales are considered to be important in stabilizing the turbulence occurring posterior to the wing. The movements of the marginal scales are closely monitored by sensory scales and bristles growing among them (A. Yoshida et al, unpublished).

Trachea and tracheoles are both important in delivering air into the wing and their patterns coincide with that of the boundary of degeneration and differentiation zones at the distal end of the wing. According to previous observations,




Figure 1. Transmission electron micrograph of the tracheole cell. Many cross-sections of the tracheoles can be observed within the cytoplasm of the tracheole cell. As the cell migrates, the tracheole is laid behind.

the pattern formation of wing epithelium is often dependent on tracheal and tracheole patterns. Basic research on the development of tracheal pattern formation is being done by scanning and transmission electron microscopy (Figure 1) to describe the exact pathway and time course of the formation of the elaborate pattern of trachea and tracheoles and to establish the cytological and developmental relationship between the formation of the tracheal pattern and the epithelial cell pattern.

II . Other research activities

This laboratory also conducts morphological observation of various animal tissues by scanning and transmission electron microscopy and immuno-electron microscopic analyses. Training in specimen preparation and instrument operation for such observations is also given. Our institute's Division of Sex Differentiation and the Laboratory of Neurophysiology were both involved in these activities.