

DIVISION OF CELL MECHANISMS



Professor
NISHIMURA, Mikio



Associate Professor
HAYASHI, Makoto

<p><i>Assistant Professors:</i></p> <p><i>Technical Staff:</i></p> <p><i>NIBB Research Fellow:</i></p> <p><i>Postdoctoral Fellow:</i></p> <p><i>Graduate Students:</i></p> <p><i>Technical Assistants:</i></p> <p><i>Secretary:</i></p>	<p><i>MANO, Shoji</i></p> <p><i>YAMADA, Kenji</i></p> <p><i>KONDO, Maki</i></p> <p><i>OIKAWA, Kazusato</i></p> <p><i>KANAI, Masatake</i></p> <p><i>GOTO, Shino</i></p> <p><i>NAKAI, Atsushi</i></p> <p><i>CUI, Songkui</i></p> <p><i>SHIBATA, Michitaro</i></p> <p><i>NISHINA, Momoko</i></p> <p><i>ARAKI, Masami</i></p> <p><i>SAITO, Miyuki</i></p> <p><i>NAKAYAMA, Tomomi</i></p> <p><i>HIKINO, Kazumi</i></p> <p><i>UEDA, Chizuru</i></p>
---	---

Since plants spread their roots in the ground, they must survive in a given environment. To adapt to the environment, they positively utilize environmental changes in the life cycle as important signals that are necessary for their survival. Plant cells can induce, degenerate and differentiate their organelles to adapt to environmental changes. The flexibility of plant organelles is the basis of the strategy for environmental adaptation in plants.

The aim of this division is to clarify the molecular mechanisms underlying the induction, differentiation, and interaction of organelles, and to understand the integrated function of individual plants through organelle dynamics. The Scientific Research on Innovative Areas “Environmental sensing of plants: Signal perception, processing and cellular responses” was started to clarify the molecular mechanisms underlying organelle differentiation and interaction.

I. Reversible transformation of plant peroxisomes

Dramatic metabolic changes that underlie the shift from heterotrophic to autotrophic growth occur in the greening of seed germination. Accompanying these metabolic changes are the functional transformations of many constitutive organelles. Etioplasts differentiate into chloroplasts while mitochondria acquire the ability to oxidize glycine. Glyoxysomes, which are peroxisomes engaged in the degradation of reserve oil via β -oxidation and the glyoxylate cycle, are transformed into leaf peroxisomes that function in several crucial steps of photorespiration. After the functional transition of glyoxysomes to leaf peroxisomes during the greening of pumpkin cotyledons, the reverse transition of leaf peroxisomes to glyoxysomes occurs during senescence. Gene expression, alternative splicing, protein translocation and protein degradation control the functional transformation between glyoxysomes and leaf peroxisomes.

II. Transcriptomics, proteomics and phenomics of plant peroxisomes

Enzymes localized in plant peroxisomes are synthesized in the cytosol and function after their post-translational transport into peroxisomes. Almost all of the peroxisomal matrix proteins contain one of two targeting signals (PTS1 and PTS2) within the molecules. PTS1 is a unique tripeptide sequence found in the carboxyl terminus of the mature proteins. In contrast, PTS2 is involved in a cleavable amino terminal presequence of peroxisomal proteins that are synthesized as a precursor protein with larger molecular mass.

We identified 256 gene candidates of PTS1- and PTS2-containing proteins and another 30 genes of non-PTS-containing proteins from the *Arabidopsis* genome. Custom-made DNA microarrays covering all these genes were used to investigate expression profiles of the peroxisomal genes in various organs. They revealed that peroxisome in root cells plays a pivotal role in polyamine catabolism. We also made a two-dimensional protein map of glyoxysomes and leaf peroxisomes isolated from *Arabidopsis* and soybean. Peptide MS fingerprinting analyses allowed us to identify novel peroxisomal membrane proteins, i.e. voltage-dependent anion-selective channel and adenine nucleotide carrier 1 (PNC1). We also found that peroxisomal membrane ATP-binding cassette transporter promotes seed germination by inducing pectin degradation under the control of abscisic acid signaling. The overall results provide us with new insights into plant peroxisomal functions.

Bioinformatic analysis of the *Arabidopsis* genome predicted the presence of 15 kinds of genes, called *PEX* genes, for peroxisomal biogenesis factors. We demonstrated that *PEX5* and *PEX7* form a cytosolic receptor complex and recognize PTS1- and PTS2-containing proteins, respectively. *PEX14* is a peroxisomal membrane docking protein that captures the receptor-cargo complex. We also comprehensively investigated whether or not these predicted *PEX* genes function in peroxisome biogenesis by generating knock-down mutants that suppress *PEX* gene expression by RNA-interference. Phenotypes of these mutants allowed us to identify the functional *PEX* genes, which can be classified into two groups: *PEX* genes regulating for peroxisomal protein import and *PEX* genes regulating for peroxisomal morphology. We continue to investigate the detailed molecular functions of other *PEX* genes. Of these, we proposed that *PEX10* is essential for the maintenance of ER morphology and for biosynthesis of cuticular wax.

III. Identification of novel components essential for peroxisome biogenesis

To better understand peroxisome biogenesis, we isolated a number of *Arabidopsis* mutants having aberrant peroxisome morphology (*apm* mutants) based on a different pattern of GFP fluorescence from the parent plant, GFP-PTS1, in which peroxisomes with normal sizes and numbers can be visualized with GFP.

Of these *apm* mutants, *APM1* gene (whose defect causes the elongation of peroxisomes and mitochondria) encodes dynamin-related protein 3A (DRP3A), one member of the dynamin family. In *apm2* and *apm4*, the GFP fluorescence is




Figure 1. Phenotype of *Arabidopsis apm* mutants. GFP fluorescence in leaf cells was observed in the parent plant, GFP-PTS1 (A), *apm1* (B), *apm3* (C) and *apm9* (D) mutants. In *apm1* and *apm3* mutants, the number of peroxisomes is dramatically decreased. However, the morphology varies in both mutants. *apm1* and *apm3* have elongated and enlarged peroxisomes, respectively. In *apm9*, GFP fluorescence is observed in the cytosol because of the efficiency of protein transport to peroxisomes. Bar indicates 20 μ m.

observed in the cytosol as well as in peroxisomes, showing the defect of protein transport to peroxisomes. We demonstrated that *APM2* and *APM4* encode proteins homologous to PEX13 and PEX12, respectively, and that *APM2/PEX13* and *APM4/PEX12* are components of the protein-translocation machinery on peroxisomal membranes. We are currently analyzing the functions of other *APM* proteins such as *APM3* and *APM9*. *apm3* and *apm9* mutants exhibit enlarged peroxisomes and defect in protein transport, respectively (Figure 1). From these analyses, we will be able to identify the components responsible for peroxisome biogenesis, and to address the mechanism at the molecular level.

IV. ER derived organelles for protein storing and defense strategy

Plant cells develop various types of endoplasmic reticulum (ER)-derived structures with specific functions. ER bodies are ER-derived compartments observed in *Arabidopsis*. They are rod-shaped structures surrounded by ribosomes, and widely distributed in the epidermal cells of whole seedlings. Undamaged rosette leaves have no ER bodies, but accumulate ER bodies after wounding or jasmonic acid treatment. This suggests that ER bodies function in the defense against herbivores. ER bodies in seedlings include *PYK10*, a β -glucosidase with an ER retention signal. *Arabidopsis nai1* mutants have no ER bodies in the entire plant and do not accumulate *PYK10*. *NAI1* encodes a transcription factor that has a basic-helix-loop-helix (bHLH) domain and regulates the expression of *PYK10* and *NAI2* (Figure 2). *Arabidopsis nai2* mutant has no ER bodies and

reduces the accumulation of *PYK10*. *NAI2* encodes a unique protein that localizes to the ER body. We found that the membrane protein of ER body 1 (*MEB1*) and *MEB2* are integral membrane proteins of the ER body. *NAI2* deficiency relocates *MEB1* and *MEB2* to the ER network. These findings indicate that *NAI2* is a key factor that enables ER body formation. Now we are investigating the function of *NAI2* on ER body formation by heterologously expressed it in onion and tobacco cells.




Figure 2. Model of ER body formation. *NAI1* regulates the expression of *NAI2* and *PYK10* in constitutive ER bodies of seedling epidermal cells. *NAI2* is responsible for the formation of ER bodies and for the accumulation of *PYK10*.

V. Vacuoles responsible for programmed cell death in plants

The vacuolar processing enzyme (VPE) belongs to the cysteine protease family found in higher plants and animals. VPE is responsible for the maturation of various types of vacuolar proteins. We revealed a novel function of VPE in various instances of programmed cell death (PCD) in plants. VPE is identified as the proteinase that exhibits caspase-1 activity in plants. The plant hypersensitive response (HR), a type of defense strategy, constitutes well-organized PCD. No HR occurs on the tobacco mosaic virus-infected leaves of VPE-deficient tobacco plants. These results suggest that VPE is involved in vacuolar collapse, which triggers PCD.

Using inhibitors for caspase-3 and the proteasome (also known to affect animal cell death), we found that the activities of both are required for bacterium-induced cell death in plants. RNA interference-mediated silencing confirmed that one of the three *Arabidopsis* proteasome catalytic subunits, *PBA1*, is required for the fusion of the vacuolar and plasma membranes, which triggers PCD.

Plants evolve a death strategy mediated by vacuolar systems, which are not seen in animals. Interestingly, vacuoles are the key players in the plant-specific cell death system.

VI. Roles of molecular chaperones on cell differentiation

Molecular chaperones are cellular proteins that function in the folding and assembly of certain other polypeptides into oligomeric structures. To clarify the roles of molecular chaperones on cell differentiation, we have purified and characterized chaperonin and HSP70s and analyzed their roles in the translocation of proteins into chloroplasts.

We found that HSP90 inhibitor induced genes with heat shock response element (HSE) motifs in their promoters, suggesting that heat shock transcription factor (HSF) is involved in the response. *Arabidopsis* HSFs interacted with HSP90.2. Thus, it appears that in the absence of heat shock, HSP90 actively suppresses HSF function. During heat shock, HSP90 is transiently inactivated, which leads to HSF activation. This data indicates that HSP90 regulates correct gene expression for heat acclimatization in plants. We also observed that HSP90 is involved in hormone responses in *Arabidopsis*. The evolutionary and functional characterizations are now being investigated.

VII. Update of The Plant Organelles Database 2 (PODB2) and release of Plant Organelles World

The Plant Organelles Database 2 (PODB2) was built to promote a comprehensive understanding of organelle dynamics. This public database is open to all researchers. PODB2 consists of four individual units: the organelles movie database, the organelle database, the functional analysis database, and external links. The organelles movie database contains time-lapse images and 3D structure rotations. The organelle database is a compilation of static image data of various tissues of several plant species at different developmental stages. The functional analysis database is a collection of protocols for plant organelle

research. The amount of included data is increasing day by day. It is expected that PODB2 will contribute to systems biology through the combination of the included data with other 'omics' data and computational analyses. In addition, we released a new website, Plant Organelles World, which is based on PODB2 as an educational tool to engage members of the non-scientific community. We expect that PODB2 and Plant Organelles World will enhance the understanding of plant organelles among researchers and the general public who want to explore plant biology.

Publication List

[Original papers]

- Kanai, M., Nishimura M., and Hayashi, M. (2010). A peroxisomal ABC transporter promotes seed germination by inducing pectin degradation under the control of *ABI5*. *Plant J.* 62, 936-947.
- Nakamura, S., Mano, S., Tanaka, Y., Ohnishi, M., Nakamori, C., Araki, M., Niwa, T., Nishimura, M., Kaminaka, H., Nakagawa, T., Sato, Y., and Ishiguro, S. (2010). Gateway binary vectors with the bialaphos resistance gene, *bar*, as a selection marker for plant transformation. *Biosci. Biotechnol. Biochem.* 74, 1315-1319.
- Shirakawa, M., Ueda, H., Shimada, T., Koumoto, Y., Shimada, T.L., Kondo, M., Takahashi, T., Okuyama, Y., Nishimura, M., and Hara-Nishimura, I. (2010). Arabidopsis Qa-SNARE SYP2 proteins localized to different subcellular regions function redundantly in vacuolar protein sorting and plant development. *Plant J.* 64, 924-935.
- Takahashi, K., Shimada, T., Kondo, M., Tamai, A., Mori, M., Nishimura, M., and Hara-Nishimura, I. (2010). Ectopic expression of an esterase, which is a candidate for the unidentified plant cutinase, causes cuticular defects in *Arabidopsis thaliana*. *Plant Cell Physiol.* 51, 123-131.

[Original paper (E-publication ahead of print)]

- Mano, S., Miwa, T., Nishikawa, S., Mimura, T., and Nishimura, M. The Plant Organelles Database 2 (PODB2): an updated resource containing movie data of plant organelle dynamics. *Plant Cell Physiol.* 2010 Nov 28.




Figure 3. The graphical user interface of the homepage in 'Plant Organelles World' (<http://podb.nibb.ac.jp/Organelle/PODBworld/en/index.html>).