

DIVISION OF MOLECULAR ENVIRONMENTAL ENDOCRINOLOGY



Professor
IGUCHI, Taisen

- Assistant Professors:** MIYAGAWA, Shinichi
 OGINO, Yukiko
- NIBB Research Fellow:** CHAKRABORTY, Tapas
- Postdoctoral Fellow:** MIYAKAWA, Hitoshi
- Technical Staff:** MIZUTANI, Takeshi
- Graduate Students:** SATO, Masaru
 TOYOTA, Kenji
 SUMIYA, Eri
 YATSU, Ryohei
 NAKAMURA, Takeshi
- Visiting Scientists:** ODA, Shigeto
 HIRAKAWA, Ikumi
 TOHYAMA, Saki
 SEBILLOT, Anthony
 WON, Seok
 LANGE, Anke
- Technical Assistants:** HAYASHI, Tomoko
 KOBAYASHI, Kaoru
- Secretary:** IMAIZUMI, Taeko

Synthetic chemicals found in the environment have the capacity to disrupt the development and function of the endocrine system in both wildlife and humans. This has drawn public concern since many of these chemicals may bind to estrogen receptors (ERs) and evoke estrogenic effects. Early evidence that exposure to estrogenic chemicals during development could pose a threat to human health came from studies of a synthetic hormone, diethylstilbestrol (DES), which was used to prevent premature birth and spontaneous abortion. Laboratory experiments showed that exposure of animals to sex hormones during critical windows of perinatal life caused irreversible alterations to the endocrine and reproductive systems of both sexes. The immune and nervous systems, bone, muscle, and the liver were also affected. Although many of these chemicals can bind to ERs in wildlife and humans, the molecular basis for the action of environmental estrogens remains poorly understood. Thus, understanding the molecular mechanisms through which environmental estrogens and sex hormones act during critical developmental windows is essential.




Figure 1. Scheme of estrogen-dependent and -independent vaginal epithelial cell proliferation in mice induced by perinatal estrogen exposure.

I. Developmental origin of adult disease: Perinatal estrogen exposure induces persistent changes in reproductive tracts

The emerging paradigm of the “embryonic/fetal origins of adult disease” provides a powerful new framework for considering the effects of endocrine disruptors on human and animal health. In 1971, prenatal DES exposure was found to result in various abnormalities of the reproductive tract in women. This syndrome was named the DES syndrome. Similar abnormalities have been demonstrated in experimental animals exposed perinatally to estrogens. Developmental estrogen exposure in mice, for example, induces persistent proliferation of vaginal epithelial cells. We found that the persistent changes in the vagina in mice exposed neonatally to estrogens result from the persistent activation of erbBs and ER α , and sustained expression of EGF-like growth factors. Currently, we are analyzing the methylation status in the mouse vagina using MeDIP (methylated DNA immunoprecipitation) coupled with a microarray (MeDIP-chip). We found several differentially methylated or demethylated DNA profiles in neonatally DES-exposed mouse vaginae and controls. We thus consider that neonatal DES exposure affects DNA methylation profiles, resulting in persistent abnormalities in mouse reproductive organs.




Figure 2. A hypothetical model for the estrogen-independent ER activation pathway in mouse vaginae.

II. Estrogen receptors of birds, reptiles, amphibians and fishes

Steroid and xenobiotic receptors (SXR) have been cloned from various animal species (fish, amphibian, reptiles, birds and mammals) by our group and we have demonstrated species-specific differences in their responses to various environmental and endogenous chemicals (receptor gene zoo). Thus, simple predictions of chemical effects based on data from a few established model species are not sufficient to develop real world risk assessments. ER and ER-like genes have been cloned from various animal species including rockshell, *Amphioxus*, lamprey, lungfish, sturgeon,

gar, roach, stickleback, mosquitofish, mangrove *Rivulus*, catshark, whale shark, Japanese giant salamander, Tokyo salamander, newt, axolotl, toad, *Silurana tropicalis*, American alligator, Nile crocodile, freshwater turtle, Japanese rat snake, Okinawa habu, and vultures. Functional studies showed that the *Amphioxus* ER sequence does not bind estrogen but *Amphioxus* steroid receptor and lamprey ER exhibited ligand-dependent transactivation, proving that invertebrate and primitive vertebrates, such as the Agnatha, have a functional ER. We found that medaka ER subtypes have their specific functions, and medaka, zebrafish and stickleback ERs are more sensitive to estrogen/estrogen-like chemical exposures than other fishes by reporter gene assay. Thus, these approaches are efficient to evaluate the relationship between species and their sensitivities to chemicals.

III. Evolutionary history and functional characterization of androgen receptor genes in jawed vertebrates

Vertebrates show diverse sexual characteristics which are regulated by androgens. To elucidate the evolutionary history and functional diversification of androgen receptor (AR) genes in vertebrates, we cloned the AR cDNAs from a shark, basal ray-finned fishes (Actinopterygii), namely bichir and sturgeon (Acipenseriformes), and teleosts including a basal teleost, arowana (Osteoglossiformes). Molecular phylogenetic analysis revealed that a gene duplication event gave rise to two different teleost ARs (α and β) and likely occurred in the actinopterygian lineage leading to teleosts after the divergence of Acipenseriformes but before the split of Osteoglossiformes. Functional analysis revealed that the shark AR activates the target gene via androgen response element by classical androgens. The teleost AR α showed unique intracellular localization with a significantly higher transactivation capacity than that of teleost AR β . These results indicate that the most ancient type of AR, as activated by the classic androgens as ligands, emerged before the Chondrichthyes-Osteichthyes split and the AR gene was duplicated during a teleost-specific gene duplication event.

IV. Gene zoo and receptor zoo

We are establishing cDNA library banks and receptor gene banks of animal species including lancelet, lamprey, sturgeon, lungfish, gar, mangrove *Rivulus*, whale shark, Japanese giant salamander, newt, *Rana rugosa*, *Silurana tropicalis*, Japanese rat snake, Okinawa habu, Florida red berry turtle, American alligator, Nile crocodile, vulture and polar bear in collaboration with the University of Pretoria, South Africa, University of Florida, Medical University of South Carolina, San Diego Zoo, USA, and the Asa Zoo in Hiroshima.

V. Sex differentiation mechanism in Daphnids

Daphnia magna has been used extensively to evaluate the organism- and population-based responses of toxicity or reproductive toxicity tests. These tests, however, provide no




Figure 3. Evolutionary relationships of androgen receptor sequences.

information about the mode of action of the tested compounds. Therefore, we applied an ecotoxicogenomic assessment of *D. magna*. We established a *Daphnia* EST database and developed an oligonucleotide-based DNA microarray with high reproducibility and demonstrated the usefulness of the array for the classification of toxic chemicals as well as for the molecular understanding of chemical toxicity in a common freshwater organism. *D. magna* reproduce asexually (parthenogenesis) when they are in an optimal environment for food, photoperiod and population density. Once environmental conditions become sub-optimal, they alter their reproductive strategy from




Figure 4. Life cycle of *Daphnia*.

asexual to sexual reproduction. Chemicals are able to affect the sex determination of *D. magna* and we found that juvenile hormone (JH) agonists (insect growth regulators), for example, induce the production of male offspring. The molecular basis of environmental sex determination is largely unknown in *D. magna*. To understand the molecular mechanisms of this phenomenon we isolated sex determination-related genes. Also, we have developed a method to inject genes into *D. magna* embryos which will allow us to study gain- and loss-of function analyses in more

detail in this species. Using these techniques, we demonstrated that DSX1 (double sex 1), one of the DM-domain genes, is essential for male differentiation in *D. magna*. To further explore the signaling cascade of sexual differentiation in *D. magna*, a gene expression profile of JH-responsive genes is essential. Thus, DNA microarray analysis has been performed in the gonads of *D. magna* exposed to fenoxycarb (synthesized JH agonist widely used as an insect growth regulator) and methyl farnesoate (JH identified in decapods) at the critical timing of JH-induced sex determination in *D. magna*. We are currently identifying JH-responsive genes in the ovary.

Publication List

[Original papers]

- Bermudez, D.S., Skotko, J.P., Ohta, Y., Boggs, A.S.P., Iguchi, T., and Guillette, L.J.Jr. (2011). Sex steroid and thyroid hormone receptor expressions in the thyroid of the American alligator (*Alligator mississippiensis*) during different life stages. *J. Morphol.* 272, 698-703.
- Chakraborty, T., Katsu, Y., Zhou, L.Y., Miyagawa, S., Nagahama, Y., and Iguchi, T. (2011). Estrogen receptors in medaka (*Oryzias latipes*) and estrogenic environmental contaminants: an *in vitro-in vivo* correlation. *J. Steroid Biochem. Mol. Biol.* 123, 115-121.
- Chakraborty, T., Shibata, Y., Zhou, L.Y., Katsu, Y., Iguchi, T., and Nagahama, Y. (2011). Differential expression of three estrogen receptor subtype mRNAs in gonads and liver from embryos to adults of the medaka, *Oryzias latipes*. *Mol. Cell. Endocrinol.* 333, 47-54.
- Kato, Y., Kobayashi, K., Watanabe, H., and Iguchi, T. (2011). Environmental sex determination in the branchiopod crustacean *Daphnia magna*: Deep conservation of a *Doublesex* gene in the sex-determining pathway. *PLoS Genetics* 7, e1001345.
- Kato, Y., Shiga, Y., Kobayashi, K., Tokishita, S., Yamagata, H., Iguchi, T., and Watanabe, H. (2011). Development of an RNA interference method in the cladoceran crustacean *Daphnia magna*. *Devel. Genes Evol.* 220, 337-345.
- Lange, A., Paull, G.C., Hamilton, P.B., Iguchi, T., and Tyler, C.R. (2011). Implications of persistent exposure to treated wastewater effluent for breeding in wild roach (*Rutilus rutilus*) populations. *Environ. Sci. Technol.* 45, 1673-1679.
- Matsumura, D., Haraguchi, R., Miyagawa, S., Motoyama, J., Nakagata, N., Meijlink, F., and Yamada, G. (2011). Genetic analysis of hedgehog signaling in ventral body wall development and the onset of omphalocele formation. *PLoS ONE* 20, e16260.
- Miyagawa, S., Matsumaru, D., Murashima, A., Omori, A., Satoh, Y., Haraguchi, R., Motoyama, J., Iguchi, T., Nakagata, N., Hui, C.C., and Yamada, G. (2011). The role of sonic hedgehog-Gli2 pathway in the masculinization of external genitalia. *Endocrinology* 152, 2894-2903.
- Moore, B.C., Milnes, M.R., Kohno, S., Katsu, Y., Iguchi, T., Woodruff, T.K., and Guillette, L.J.Jr. (2011). Altered gonadal expression of TGF- β superfamily signaling factors in environmental contaminant-exposed juvenile alligators. *J. Steroid Biochem. Mol. Biol.* 127, 58-63.
- Murashima, A., Miyagawa, S., Ogino, Y., Nishida-Fukuda, H., Araki, K., Matsumoto, T., Kaneko, T., Yoshinaga, K., Yamamura, K.I., Kurita, T., Kato, S., Moon, A.M., and Yamada, G. (2011). Essential roles of androgen signaling in Wolffian duct stabilization and epididymal cell differentiation. *Endocrinology* 152, 1640-1651.
- Nakajima, T., Hayashi, S., Iguchi, T., and Sato, T. (2011). The role of fibroblast growth factors on the differentiation of vaginal epithelium of neonatal mice. *Differentiation* 82, 28-37.
- Nakajima, T., Iguchi, T., and Sato, T. (2011). Involvement of activin signaling in abnormalities of the mouse vagina exposed neonatally to diethylstilbestrol. *Cell Tiss. Res.* 344, 527-538.
- Southam, A.D., Lange, A., Hines, A., Hill, E.M., Katsu, Y., Iguchi, T., Tyler, C.R., and Viant, M.R. (2011). Metabolomics reveals target and off-target toxicities of a model organophosphate pesticide to roach (*Rutilus rutilus*): Implications for biomonitoring. *Environ. Sci. Technol.* 45, 3759-3767.
- Urushitani, H., Katsu, Y., Miyagawa, S., Kohno, S., Ohta, Y., Guillette, L.J.Jr., and Iguchi, T. (2011). Molecular cloning of anti-Müllerian hormone from the American alligator, *Alligator mississippiensis*. *Mol. Cell. Endocrinol.* 333, 190-199.
- Urushitani, H., Katsu, Y., Ohta, Y., Shiraishi, H., Iguchi, T., and Horiguchi, T. (2011). Cloning and characterization of retinoid X receptor (RXR) isoforms in the rock shell, *Thais clavigera*. *Aquat. Toxicol.* 103, 101-111.

[Review articles]

- Blumberg, B., Iguchi, T., and Odermatt, A. (2011). Editorial: Endocrine disrupting chemicals. *J. Steroid Biochem. Mol. Biol.* 127, 1-3.
- Celander, M.C., Goldstone, J.V., Denslow, N.D., Iguchi, T., Kille, P., Meyerhoff, R.D., Smith, B.A., Hutchinson, T.H., and Wheeler, J.R. (2011). Species extrapolation for the 21st century. *Environ. Toxicol. Chem.* 30, 52-63.
- Hayes, T.B., Beasley, V.R., de Solla, S., Iguchi, T., Ingraham, H., Kestemont, P., Kniewald, J., Kniewald, Z., Langlois, V.S., Luque, E.H., McCoy, K.A., Muñoz-de-Toro, M., Oka, T., Oliveira, C.A., Orton, F., Ruby, S., Suzawa, M., Tavera-Mendoza, L.E., Trudeau, V.L., Victor-Costa, A.B., and Willingham, E. (2011). Demasculinization and feminization of male gonads by atrazine: consistent effects across vertebrate classes. *J. Steroid Biochem. Mol. Biol.* 127, 64-73.
- Miyagawa, S., Sato, M., and Iguchi, T. (2011). Molecular mechanisms of induction of persistent changes by estrogenic chemicals on female reproductive tracts and external genitalia. *J. Steroid Biochem. Mol. Biol.* 127, 51-57.
- Ogino, Y., Miyagawa, S., Katoh, H., Prins, G.S., Iguchi, T., and Yamada, G. (2011). Essential functions of androgen signaling emerged through the developmental analysis of vertebrate sex characteristics. *Evol. Devel.* 13, 315-325.
- Watanabe, H., Kato, Y., and Iguchi, T. (2011). Application of ecotoxicogenomics for understanding mode of action of chemicals and species extrapolation. In: *Handbook of Systems Toxicology*. Eds. Casciano, D.A. and Sahu, S.C., John Wiley & Sons, Ltd. pp.67-72.