## LABORATORY FOR BIOTHERMOLOGY



Specially Appointed Associate Professor KAMEI, Yasuhiro

NIBB Research Fellow: CREST Researcher: JSPS Postdoctral Fellow: Technical Assistant: SAKAMOTO, Joe KAMIKAWA, Yuko SUZUKI, Miyuki KINOSHITA, Chie TAMADA, Tomoko

Temperature is an important parameter for living organisms. Cell activity is affected by temperature since the reaction ratio and stability of molecules in cells depends on it. Environmental temperatures which organisms function in are variable, and the biological system that regulates body temperature, homeostasis, differs among various organisms. Furthermore, temperature in cells or at the molecular level has not been widely discussed to date. Nano-scale thermometers utilizing fluorescent proteins and dyes have been developed by numerous research groups including ours, and heterogeneity of temperatures in the micro environments of living cells have been reported on by said groups (Okabe et al, Nat. Commun. 2014, Kiyonaka et al, Nat. Methods 2014, and Nakano et al, PLoS One 2017). However, the meaning and mechanisms in single cells remain unclear. Thus, our group is embarking upon a new research field, biothermology, through the investigation of the nature of temperature in living organisms ranging from nano to macro.



Figure 1. Schematic illustration of heat shock response of cells (left) and an infrared laser-evoked gene operator (IR-LEGO) microscope system.

Our research group employs various and original biothermological research technologies . One of these is the infrared (IR) laser application, which enables single-cell or subcellular local heating by focusing IR through microscope objective. This technology was originally developed for laser induced gene expression in the targeting of single-cells through the heat shock response, called IR-LEGO; InfraRed Laser Evoked Gene Operator (Kamei et al, Nat. Methods 2009) (Figure 1). This local heating technique can be used to analyze the thermodynamics of a cell via the use of a nano-scale thermometer; the second of our abovementioned original techniques. To analyze temperature distribution at subcellular resolutions, a nano-scale thermometer is required. For this reason, we have focused on fluorescent proteins, and developed a genetically encoded thermometer, the so called gTEMP, with Dr. Takeharu Nagai from Osaka University (Nakano et al, PLoS One 2017). This thermometer has some unique properties such as rapid response and applicability to wide temperature range. By utilizing this thermometer, we are developing a high-speed thermal imaging microscope system combined with IR irradiation optics. Through the analysis of thermal dynamics in cells, we are trying to reveal how temperature heterogeneity is generated within a single cell and to consider its meaning from a biological standpoint.

Furthermore, our group is trying to improve the IR-LEGO technique itself. We initially applied this system to many organisms including animals and plants, such as medaka, nematode, flies, frogs, and Araidopsis, moss, in collaborative research projects in cooperation with other laboratories. Because the heat shock response (HSR) is a transient response, gene expressions after IR-LEGO operation are also transient. Despite this, some of our collaborators wanted to realize cell fate mapping, which requires a long-term gene expression after IR irradiation. Accordingly, we employed Cre/LoxP recombination system. As shown in Figure 2, we established a cre driver line, which possesses a heat shock promoter and loxP effector line which can permanently label irradiated cell lineages by fluorescent proteins. By using these transgenic lines, evidence was provided that indicated that exoskeletal tissues in trunk region came from the mesoderm (Shimada et al, Nat. Commun. 2013).



Figure 2. Long-term gene expression with cre/loxP recombination system and an example of a practical experiment in medaka ranging embryos to adults.

The HSR is a conservative stress response system found in almost all organisms. In contrast, organisms live in various temperatures, and HSR essentially is the upper limit of habitable temperature range for each organism. This means that the preset temperature of HSR may differ among organisms. A key factor of this limitation is heat shock factor 1 (HSF1); a homologue within each organism. By substituting HSF1, it may be possible to control the preset temperature of HSR, and this substitution could be applied to the improvement of HSR efficiency in IR-LEGO technology. We are now trying to lower the HSR temperature of medaka by substituting Japanese flounder HSF1. This HSF1 substitution project provides us with various significant information and evidence related to molecular evolution and adaptation to environmental temperature. Therefore, this project also contributes to the study of biothermology.

We also promote other collaborative studies using microscopic techniques. One is a "clear observation project" for deep seeing into living organisms using adaptive optics (AO), which have been thoroughly developed in the field of astronomy as a key technology found in large telescopes, such as the Subaru telescope in Hawaii. Although observation using Earth based telescopes may be interfered with due to fluctuations in the atmosphere, AO can mitigate this. However, living materials have particular refractive indexes, so some organelles may hinder the ideal optical path for microscope observation, which is similar to the situation that exists regarding the atmosphere and telescopes. AO can also compensate for this disturbance by sensing and correcting wave fronts by using a wave front sensor and deformable mirror. Hence, we have developed a custom-made wide-field microscope equipped with an AO system for the observation of living organisms in collaboration with Dr. Yosuke Tamada from Utsunomiya University and Dr. Yutaka Hayano from the National Astronomical Observatory of Japan (NAOJ). By utilizing this microscopic system, we have successfully acquired high-resolution bright field and fluorescent images of living cells. Furthermore, AO can also be applied to IR-LEGO technology to improve IR energy focusing of irradiation in a manner similar to improving the sight of living organisms.

## **Publication List:**

[Original papers]

- Abe, E., Yasugi, M., Takeuchi, H., Watanabe, E., Kamei, Y., and Yamamoto, H. (2019). Development of omnidirectional aerial display with aerial imaging by retro-reflection (AIRR) for behavioral biology experiments. Opt. Rev. 26, 221-229. doi: 10.1007/s10043-019-00502-w
- Fujikawa, Y., Ishikawa-Fujiwara, T., Kuo, T., Shinkai, N., Shogi, T., Kawasaki, T., Kamei, Y., Sakuraba, Y., Sato, A., Kinoshita, M., Gondo, Y., Yuba, S., Tsujimura, T., Sese, J., and Todo, T. (2020). Involvement of Rev1 in alkylating agent-induced loss of heterozygosity in *Oryzias latipes*. Genes Cells 25, 124-138. doi: 10.1111/gtc.12746
- Furukawa, F., Hamasaki, S., Hara, S., Uchimura, T., Shiraishi, E., Osafune, N., Takagi, H., Yazawa, T., Kamei, Y., and Kitano, T. (2019). Heat shock factor 1 protects germ cell proliferation during early ovarian differentiation in medaka. Sci. Rep. 9, 6927. doi: 10.1038/s41598-019-43472-4
- Harada, Y., Matsuo, M., Kamei, Y., Goto, M., and Fukamachi, S. (2019). Evolutionary history of the medaka long-wavelength sensitive genes and effects of artificial regression by gene loss on behavioural photosensitivity. Sci. Rep. 9, 2726. doi: 10.1038/s41598-019-39978-6
- Hwang, D., Wada, S., Takahashi, A., Urawa, H., Kamei, Y., and Nishikawa, S. (2019). Development of a heat-inducible gene expression system using female gametophytes of *Arabidopsis thaliana*. Plant Cell Physiol. 60, 2564-2572. 10.1093/pcp/pcz148
- Ikehata, H., Mori, T., Kamei, Y., Douki, T., Cadet, J., and Yamamoto, M. (2020). Wavelength- and tissue-dependent variations in the mutagenicity of cyclobutane pyrimidine dimers in mouse skin. Photochem. Photobiol. 96, 94-104. doi: 10.1111/php.13159
- Kosugi, M., Ozawa, S., Takahashi, Y., Kamei, Y., Itoh, S., Kudoh, S., Kashino, Y., and Koike, H. (2020). Red-shifted chlorophyll a bands allow uphill energy transfer to photosystem II reaction centers in an aerial green alga, *Prasiola crispa*, harvested in Antarctica. BBA Bioenergetics *1861*, 148139. doi: 10.1016/j.bbabio.2019.148139
- Nakane, Y., Shinomiya, A., Ota, W., Ikegami, K., Shimmura, T., Higashi, S., Kamei, Y., and Yoshimura, T. (2019). Action spectrum for photoperiodic control of thyroid-stimulating hormone in Japanese quail (*Coturnix japonica*). PLoS One *14*, e022106. doi: 10.1371/journal. pone.0222106

- Nakayama, T., Shimmura, T., Shinomiya, A., Okimura, K., Takehana, Y., Furukawa, Y., Shimo, T., Senga, T., Nakatsukasa, M., Nishimura, T., Tanaka, M., Okubo, K., Kamei, Y., Naruse, K., and Yoshimura, T. (2019). Seasonal regulation of the lncRNA LDAIR modulates self-protective behaviours during the breeding season. Nat. Ecol. Evol. *3*, 845-852. doi: 10.1038/s41559-019-0866-6
- Shikata, T., Takahashi, F., Nishide, H., Shigenobu, S., Kamei, Y., Sakamoto, S., Yuasa, K., Nishiyama, Y., Yamasaki, Y., and Uchiyama, I. (2019). RNA-Seq analysis reveals genes related to photoreception, nutrient uptake, and toxicity in a noxious red-tide raphidophyte *Chattonella antiqua*. Front. Microbiol. 10, 1764. doi: 10.3389/ fmicb.2019.01764
- Uchimura, T., Hara, S., Yazawa, T., Kamei, Y., and Kitano, T. (2019). Involvement of heat shock proteins on the transcriptional regulation of corticotropin-releasing hormone in medaka. Front. Endocrinol. 10, 529. doi: 10.3389/fendo.2019.00529
- Yokoi, S., Naruse, K., Kamei, Y., Ansai, S., Kinoshita, M., Mito, M., Iwasaki, S., Inoue, S., Okuyama, T., Nakagawa, S., Young, L.J., and Takeuchi, H. (2020). Sexually dimorphic role of oxytocin in medaka mate choice. Proc. Natl. Acad. Sci. USA *117*, 4802-4808. doi: 10.1073/ pnas.1921446117

## [Review articles]

- Abe, G., Hayashi, T., Yoshida, K., Yoshida, T., Kudoh, H., Sakamoto, J., Konishi, A., Kamei, Y., Takeuchi, T., Tamura, K., and Yokoyama, H. (2020). Insights regarding skin regeneration in non-amniote vertebrates: Skin regeneration without scar formation and potential step-up to a higher level of regeneration. Semin. Cell Dev. Biol. 100, 109-121. doi: 10.1016/j.semcdb.2019.11.014
- Murata, K., Kinoshita, M., Naruse, K., Tanaka, M., and Kamei, Y. eds. (2020). Medaka: Biology, Managements, and Experimental Protocols, Volume 2. (Wiley-Blackwell, Iowa, USA). ISBN: 978-1-119-57529-0