

**LABORATORY FOR SPATIOTEMPORAL REGULATIONS**



Associate Professor  
**NONAKA, Shigenori**

Specially Appointed Assistant Professor:


*YOKE, Hiroshi*

Postdoctoral Fellow: *TANIGUCHI, Atsushi*

Technical Assistant: *ISHIBASHI, Tomoko*



Our laboratory is currently pursuing two paths of scientific inquiry that are of interest to developmental biology. One is to clarify the mechanism of left-right (L-R) determination. The other is to develop imaging technologies for biological analyses.



Visual overview of this lab's work.

**I. Initial step for left-right asymmetry**

In mammalian development, initial left-right asymmetry first appears as cilia-driven leftward fluid flow on the ventral surface of the node, and this flow provides the initial cue for asymmetric gene expressions that actually determine asymmetric morphogenesis. The mechanisms that convert the flow to asymmetric gene expression, *i.e.* the flow sensing mechanism, remain controversial, with several models being proposed, and the involvement of  $Ca^{2+}$  being suggested.




Figure 1. Left: Distribution of  $Ca^{2+}$  elevation in a 2-somite wild-type node. Right: Time course of  $Ca^{2+}$  elevation frequency at the left and the right sides.

We pursued this question by measuring  $Ca^{2+}$  dynamics in the node and found that the node cells apparently cause stochastic elevation of  $Ca^{2+}$ . The spatiotemporal distribution is equal on the left and right sides, but becomes more prevalent on the left after the late headfold stage, when flow-induced commitment occurs. This asymmetry was disrupted in dynein mutant *iv/iv* and *pkd2<sup>-/-</sup>* mutants, in accordance to their left-right phenotypes.

Based on these findings, we are now examining the validity of the two major proposed models of flow sensing, mechanosensing and chemosensing, as well as a third novel mechanism.

**II. Development of light-sheet microscopy**

Light-sheet microscopy has become popular during this decade due to benefits such as low photodamage and fast image acquisition. We are now running two light-sheet microscopes, one commercial and the other self-made, and are maintaining them for collaborations and our own research interest (this being left-right asymmetry).

Over several years, we have developed a fast light-sheet microscope named ezDSLIM, in which the excitation light sheet and detection objective move instead of the specimen for z-scanning. We installed an electrically tunable lens (ETL) to achieve greater speed and the exclusion of mechanical vibrations that disturb continuous 4D imaging of floating or loosely held specimens.

Our light-sheet microscopes are available to other researchers via NIBB's Collaborative Research and MEXT's Advanced Bioimaging Support (ABiS) programs. Numerous collaborations are in progress including observation of moving cell migration in zebrafish embryos, cleared mouse brains, etc.




Figure 2. Images of floating volvox taken by ezDSLIM with ETL. Left: Single optical section. Right: Maximum intensity projection.

**Publication List:**

[Original papers]

- Hirata, Y., Matsuo, M., Kurihara, K., Suzuki, K., Nonaka, S., and Sugawara, T. (2021). Colocalization Analysis of Lipo-Deoxyribozyme Consisting of DNA and Protic Catalysts in a Vesicle-Based Proto-cellular Membrane Investigated by Confocal Microscopy. *Life (Basel)* 11, 1364. DOI: 10.3390/life11121364
- Ohmura, T., Nishigami, Y., Taniguchi, A., Nonaka, S., Ishikawa, T., and Ichikawa, M. (2021). Near-wall rheotaxis of the ciliate *Tetrahymena* induced by the kinesthetic sensing of cilia. *Sci. Adv.* 7, eabi5878. DOI: 10.1126/sciadv.abi5878